

Quantifying Model Uncertainties in the Space of Probability Measures

Jinqiao Duan¹, Ting Gao¹ and Guowei He² *

*1. Institute for Pure and Applied Mathematics
University of California, Los Angeles, CA 90095, USA*

&

*Department of Applied Mathematics, Illinois Institute of Technology
Chicago, IL 60616, USA*

E-mail: duan@iit.edu tgao5@iit.edu

*2. Laboratory for Nonlinear Mechanics
Institute of Mechanics, Chinese Academy of Sciences
Beijing 100080, China*

E-mail: hgw@lnm.imech.ac.cn

Due to lack of scientific understanding, some mechanisms may be missing in mathematical modeling of complex phenomena in science and engineering. These mathematical models thus contain some uncertainties such as uncertain parameters. One method to estimate these parameters is based on pathwise observations, i.e., quantifying model uncertainty in the space of sample paths for system evolution. Another method is devised here to estimate uncertain parameters, or unknown system functions, based on experimental observations of probability distributions for system evolution. This is called the quantification of model uncertainties in the space of probability measures. A few examples are presented to demonstrate this method, analytically or numerically.

Keywords: Model uncertainty; Stochastic differential equations (SDEs); Probability measures; Hellinger distance; Stationary probability density; Parameter estimation; Numerical simulation

2000 AMS Subject Classification : 60H40

1 . Introduction

In this chapter we discuss some issues about quantification of model uncertainties in complex dynamical systems.

*Partly supported by the NSF grant 1025422, NSFC grant 10971225, the open funding from the Laboratory for Nonlinear Mechanics at the Chinese Academy of Sciences, and the Fundamental Research Funds for the Central Universities, HUST 2010ZD037.

Mathematical models for scientific and engineering systems often involve with some uncertainties. We may roughly classify such uncertainties into two kinds. The first kind of uncertainties may be called *model uncertainty*. They are due to physical processes that are not well understood or not well-observed, and thus are not or not well represented in the mathematical models.

The second kind of uncertainties may be called *simulation uncertainty*. This arises in numerical simulations of multiscale systems that display a wide range of spatial and temporal scales, with no clear scale separation. Due to the limitations of computer power, not all scales of variability can be explicitly simulated or resolved.

These uncertainties are sometimes also called *unresolved scales*, as they are not represented (i.e., not *resolved*) in modeling or simulation. Although these unresolved scales may be very small or very fast, their long time impact on the resolved simulation may be delicate (i.e., may be negligible or may have significant effects,^{12,60} or in other words, uncertain). Thus, to take the effects of unresolved scales on the resolved scales into account, representations or parameterizations of these effects are desirable.

Model uncertainties have been considered in, for example,^{19,26,41,44–46} Research works relevant for parameterizing unresolved scales include,^{6–8,18,25,27,31,39,50,57,58} among others. Stochastically representing unresolved scales in fluid dynamics has considered as well.^{35,40,52}

In this chapter, we only consider model uncertainties. Specifically, we consider dynamical systems containing uncertain parameters or unknown system functions, and examine how to estimate these parameters, using observed probability distributions of the system evolution.

After briefly comment on estimating uncertain parameters based on observed sample paths for the system evolution in §2, we then, in §3, propose a method of estimating uncertain parameters based on observed probability distributions (i.e., probability measures) and present a few examples to demonstrate this method, analytically or numerically.

2 . Quantifying uncertainty in the space of paths

Since random fluctuations are common in the real world, mathematical models for complex systems are often subject to uncertainties, such as fluctuating forces, uncertain parameters, or random boundary conditions.^{19,26,41,53,56,59} Stochastic differential equations (SDEs) such as

$$dX = b(X)dt + \sigma(X)dB_t, \quad (2.1)$$

are appropriate models for many of these systems.^{5,49,56} Here B_t is a Brownian motion or Wiener process, the drift $b(X)$ and diffusion $\sigma(X)$ contain uncertain parameters (or $b(\cdot)$ and $\sigma(\cdot)$ are unknown), to be estimated based on observations.

For example, the Langevin type models are stochastic differential equations de-

scribing various phenomena in physics, biology, and other fields. SDEs are used to model various price processes, exchange rates, and interest rates, among others, in finance. Noises in these SDEs may be modeled as a generalized time derivative of some distinguished stochastic processes, such as Brownian motion (BM) or other processes.

We are interested in estimating parameters contained in the stochastic differential equation (2.1), so that we obtain computational models useful for investigating complex dynamics under uncertainty.

Theoretical results on parameter estimations for SDEs driven by Brownian motion are relatively well developed,^{4,9,13,23,28,34,47} and various numerical simulations for these parameter estimations^{1,3,42,47} are also implemented. See⁶¹ for a more recent review about estimating and computing uncertain parameters, when dynamical systems are submit to colored or non-Gaussian noises.

These research works on estimating uncertain parameters in dynamical systems are based on observations of sample paths. In the next section, we devise a method to estimate uncertain parameters based on observations of probability distributions of the system evolution.

3 . Quantifying uncertainty in the space of probability measures

Consider a dynamical system with model uncertainty, modeled by a scalar SDE

$$dX = b(X)dt + \sigma(X)dB_t, \quad X(0) = x_0, \quad (3.1)$$

where the drift $b(X)$ and diffusion $\sigma(X)$ contain uncertain parameters, to be estimated based on observations of probability distributions (i.e., probability measures) of the system paths X_t .

To this end, we need to introduce the Hellinger distance¹⁰ between two probability measures. It is used to quantify the similarity between two probability distributions. This is a metric in the space of probability measures.

For our purpose here, we define the Hellinger distance $H(f, g)$ between two probability density functions $p(x)$ and $q(x)$ as follows

$$H^2(p, q) \triangleq \frac{1}{2} \int_{\mathbb{R}^1} (\sqrt{p(x)} - \sqrt{q(x)})^2 dx. \quad (3.2)$$

The Hellinger distance $H(p, q)$ satisfies the property: $0 \leq H(p, q) \leq 1$.

We estimate uncertain parameters by minimizing the Hellinger distance between the true probability density p for the solution process $X(t)$ and its observed probability density q . In reality, the probability density p has to be numerically formulated or discretized. But in order to demonstrate the method, we consider two examples for which the true probability density p can be analytically formulated. In the first example, we minimize the Hellinger distance between the true stationary probability density for the solution process $X(t)$ and its observed stationary probability

density, while in the second example, we do this for time-dependent probability densities.

3.1. Observation of stationary probability distributions

Under appropriate conditions on b and σ (see,³² p.170), such as, $b \leq 0$ and $\sigma \neq 0$ as well as some smoothness requirements, there exists a stationary probability density $p(x)$ for the SDE (3.1), as a solution of the steady Fokker-Planck equation $p_{xx} + (b \sin(x)p)_x = 0$,

$$p(x) = \frac{C}{\sigma^2(x)} e^{\int_{x^*}^x \frac{2b(y)}{\sigma^2(y)} dy},$$

where the positive normalization constant C is chosen so that $p \geq 0$ and $\int_{\mathbb{R}} p(x) dx = 1$, i.e.,

$$C \triangleq 1 / \int_{-\infty}^{\infty} \frac{e^{\int_{x^*}^x \frac{2b(y)}{\sigma^2(y)} dy}}{\sigma^2(x)} dx.$$

Note that x^* here may be an arbitrary point so that the integral $\int_{x^*}^x \frac{2b(y)}{\sigma^2(y)} dy$ exists (say, take $x^* = 0$ if that is possible).

Example 3.1. (i) A special case: Langevin equation

$$dX = -bXdt + dB_t,$$

with parameter $b > 0$. Given an “observation” of the stationary probability density $q(x) = \frac{1}{\sqrt{\pi}} e^{-x^2}$. Find a b so that the Hellinger distance $F(b) = \frac{1}{2} \int_{\mathbb{R}} [\sqrt{p(x)} - \sqrt{q(x)}]^2 dx$ is minimized.

(ii) A more general case:

$$dX = b(X)dt + dB_t,$$

with function $b(x) \leq 0$. Given an “observation” of the stationary probability density $q(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ (the Cauchy distribution). Find a function $b(x) \leq 0$ so that the Hellinger distance $F(b(x)) = \frac{1}{2} \int_{\mathbb{R}} [\sqrt{p(x)} - \sqrt{q(x)}]^2 dx$ is minimized.

Solution:

(i) The true stationary probability density for the solution process X_t is

$$p(x) = \frac{\sqrt{b}}{\sqrt{\pi}} e^{-bx^2}.$$

Insert p, q into the Hellinger distance $F(b)$, which is now an algebraic function of parameter $b > 0$. Thus we use deterministic calculus to find a minimizer b (possibly by hand, or Matlab if needed). Note: $\int_{\mathbb{R}} e^{-z^2} dz = \sqrt{\pi}$.

To minimize the Hellinger distance $F(b)$, we calculate its derivative

$$\begin{aligned} F'(b) &= \frac{1}{2} \int_{\mathbb{R}} e^{-bx^2} \left(\frac{1}{2\sqrt{\pi b}} - \frac{\sqrt{b}}{\sqrt{\pi}} x^2 \right) dx - \frac{1}{\sqrt{\pi}} \int_{\mathbb{R}} e^{-\frac{(1+b)x^2}{2}} \left(\frac{1}{4} b^{-\frac{3}{4}} - \frac{x^2}{2} b^{\frac{1}{4}} \right) dx \\ &= \frac{1}{\sqrt{2}} b^{\frac{1}{4}} (1+b)^{-\frac{3}{2}} - \frac{1}{2\sqrt{2}} b^{-\frac{3}{4}} (1+b)^{-\frac{1}{2}} = 0. \end{aligned}$$

Therefore,

$$b^{\frac{1}{4}} (1+b)^{-\frac{3}{2}} = \frac{b^{-\frac{3}{4}} (1+b)^{-\frac{1}{2}}}{2}.$$

Thus we obtain the parameter $b = 1$.

(ii) The true stationary probability density for the solution process X_t is

$$p(x) = \frac{e^{2 \int_0^x b(y) dy}}{\int_{-\infty}^{\infty} e^{2 \int_0^x b(y) dy} dx}.$$

Insert p, q into the Hellinger distance $F(b(x))$, which is now a functional of $b(x)$ and thus we use calculus of variations (on $F(b(x))$) to find a minimizer $b(x)$. We then derive the Euler-Lagrange equation to be satisfied by $b(x)$, together with appropriate boundary conditions (needed for $p(x) \geq 0$ and $\int_{\mathbb{R}} p(x) dx = 1$).

To this end, we calculate, for an arbitrary “variations” $h(x)$

$$\begin{aligned} &F(b(x) + \varepsilon h(x)) \\ &= \frac{1}{2} \int_{\mathbb{R}} \left[\frac{e^{2 \int_0^x (b(y) + \varepsilon h(y)) dy}}{\int_{-\infty}^{\infty} e^{2 \int_0^x (b(y) + \varepsilon h(y)) dy} dx} - \frac{2e^{\int_0^x (b(y) + \varepsilon h(y)) dx}}{\sqrt{\pi} \sqrt{1+x^2} \sqrt{\int_{-\infty}^{\infty} e^{2 \int_0^x (b(y) + \varepsilon h(y)) dy} dx}} + \frac{1}{\pi(1+x^2)} \right] dx. \end{aligned}$$

The Euler-Lagrange equation for $b(x)$ comes from: $\frac{d}{d\varepsilon} \big|_{\varepsilon=0} F(b(x) + \varepsilon h(x)) = 0$ for arbitrary “variations” $h(x)$. In fact, the Euler-Lagrange equation for $b(x)$ is

$$\int_{\mathbb{R}} e^{\int_0^x b(w) dw} \left(e^{\int_0^x b(w) dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w) dw} dx}}{\sqrt{\pi(1+x^2)}} \right) \cdot \int_{\mathbb{R}} \left(e^{2 \int_0^z b(w) dw} \int_z^x h(y) dy \right) dz = 0.$$

After changing the order of integration (first on y and z then on y and x), we have

$$\left\{ \int_y^{\infty} e^{\int_0^x b(w) dw} \left(e^{\int_0^x b(w) dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w) dw} dx}}{\sqrt{\pi(1+x^2)}} \right) dx \cdot \int_{-\infty}^y e^{2 \int_0^z b(w) dw} dz \right\} h(y) dy = 0$$

holds for all $h(y)$.

Therefore,

$$\int_y^{\infty} e^{\int_0^x b(w) dw} \left(e^{\int_0^x b(w) dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w) dw} dx}}{\sqrt{\pi(1+x^2)}} \right) dx \cdot \int_{-\infty}^y e^{2 \int_0^z b(w) dw} dz = 0.$$

Since $\int_{-\infty}^y e^{2 \int_0^x b(w)dw} dz > 0$, we further obtain

$$\int_y^\infty e^{\int_0^x b(w)dw} \left(e^{\int_0^x b(w)dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx}}{\sqrt{\pi(1+x^2)}} \right) dx = 0$$

for $y \in \mathbb{R}$. Then taking the derivative of the above equation with respect to y , we arrive at

$$e^{\int_0^y b(w)dw} = \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx}}{\sqrt{\pi(1+y^2)}}, \quad \forall y \in \mathbb{R}^1.$$

Thus, after taking the ‘square’ and ‘ln’ on both sides of the above equation, we get

$$2 \int_0^y b(w)dw = \ln \left(\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx \right) - \ln(\pi(1+y^2)).$$

Finally taking the derivative with respect to y , we have

$$b(y) = -\frac{y}{\pi(1+y^2)} \quad y \in \mathbb{R}^1.$$

Also note that we only need $\int_0^y b(w)dw < 0$ for all $y \in \mathbb{R}^1$ for the stationary probability density to make sense.

3.2. Observation of time-dependent probability distributions

Consider a scalar SDE

$$dX = b(X)dt + \sigma(X)dB_t, \quad X(0) = x_0. \quad (3.3)$$

The Fokker-Planck equation^{20,43,54} for the probability density $p(x, t) \triangleq p(x, t; x_0, 0)$ for the solution $X(t, x_0)$ is

$$p_t = \frac{1}{2}(\sigma^2(x)p(x, t))_{xx} - (b(x)p(x, t))_x, \quad p(x, 0) = \delta(x_0). \quad (3.4)$$

With an observation of $p(x, t)$, we can estimate parameters, or $b(\cdot)$, or $\sigma(\cdot)$, by examining the inverse problem of the Fokker-Planck equation (3.4). For more information about inverse problems of partial differential equations, see.²⁹

Let us look at a specific example.

Example 3.2. Consider a scalar SDE

$$dX = -b \sin(X)dt + \sqrt{2} dB_t, \quad X(0) = 0. \quad (3.5)$$

(i) Assume that an observation obtained for p to be

$$q_1(x, t) = \frac{1}{2\sqrt{\pi t}} e^{-\frac{x^2}{4t}}. \quad (3.6)$$

Find the parameter b by minimizing the Hellinger distance $H(p, q_1)$.

(ii) Assume that another observation obtained for p to be

$$q_2(x, t) = \frac{\sqrt{t}}{\pi(t + x^2)}. \quad (3.7)$$

Find the parameter b by minimizing the Hellinger distance $H(p, q_2)$.

Solution:

The Fokker-Planck equation for (3.5) is

$$p_t = p_{xx} + (b \sin(x)p)_x, \quad p(x, 0) = \delta(0). \quad (3.8)$$

In this case, we define the Hellinger distance:

$$H_i^2(b) = \max_{t \in [0, T]} \int_{-\infty}^{\infty} (\sqrt{q_i(x, t)} - \sqrt{p(x, t, b)})^2 dx$$

where $i = 1, 2$, and T is the time period when $q_i(x, t)$ are observed. We numerical find b by minimizing $H_i(b)$.

The observation $q_1(x, t)$ is plotted in Figure 0.1.




Fig. 0.1. Observation $q_1(x, t)$: $t = 5$ (top) and $t = 30$ (bottom).

By the definition $H_1^2(b) = \max_{t \in [0, T]} \int_{-\infty}^{\infty} (\sqrt{q_1(x, t)} - \sqrt{p(x, t, b)})^2 dx$, we have the plot of $H_1(b)$ in Figure 0.2. And whatever T is, $H_1(b)$ is always minimized when $b = 0$. This gives us the parameter value $b = 0$.

The observation $q_2(x, t)$ is plotted in Figure 0.3.

By the definition $H_2^2(b) = \max_{t \in [0, T]} \int_{-\infty}^{\infty} (\sqrt{q_2(x, t)} - \sqrt{p(x, t, b)})^2 dx$, we have the plot of $H_2(b)$ in Figure 0.4. So we see that if $T = 5$, $H_2(b)$ is minimized when $b = 0.7$ and if $T = 30$, $H_2(b)$ is minimized when $b = 0.6$.

Acknowledgements. We would like to thank Huijie Qiao and Xiangjun Wang for helpful discussions.

References

1. Y. Aït-Sahalia, Maximum-likelihood estimation of discretely-sampled diffusions: a closed-form approximation approach, *Econometrica* **70** (2002), 223-262.
2. Y. Aït-Sahalia and P. A. Mykland, Estimators of diffusions with randomly spaced




Fig. 0.2. Squared Hellinger distance between $p(x, t)$ and the observation $q_1(x, t)$: $T = 5$ (top) and $T = 30$ (bottom).

- discrete observations: a general theory, *The Annals of Statistics* **32**(5)(2004), 2186-2222.
3. Y. Aït-Sahalia and P. A. Mykland, The effects of random and discrete sampling when estimating continuous-time diffusions, *Econometrica* **71**(2)(2003), 483-549.
 4. S. Alizadeh, M. W. Brandt and F. X. Diebold, Range-based estimation of stochastic volatility models, *The Journal of Finance* **57**(3)(2002), 1047-1091.
 5. L. Arnold. *Random Dynamical Systems*. Springer, New York, 1998.
 6. L. Arnold, Hasselmann's program revisited: The analysis of stochasticity in deterministic climate models. In J.-S. von Storch and P. Imkeller, editors, *Stochastic climate models*. pages 141-158, Boston, 2001. Birkhäuser.
 7. P. S. Berloff, Random-forcing model of the mesoscale oceanic eddies. *J. Fluid Mech.* **529** (2005), 71-95.
 8. L.C. Berselli, T. Iliescu and W. J. Layton. *Mathematics of Large Eddy Simulation of Turbulent Flows*. Springer Verlag, 2005.
 9. J. P. N. Bishwal, *Parameter Estimation in Stochastic Differential Equations*, Springer, New York, 2007.
 10. S. Cha, Comprehensive Survey on Distance/Similarity Measures between Probability Density Functions. *International Journal of Mathematical Models and Methods in Applied Sciences*, vol. 1(4), 2007 pp.300-307.




Fig. 0.3. Observation $q_2(x, t)$: $t = 5$ (top) and $t = 30$ (bottom).

11. B. Chen and J. Duan, Stochastic Quantification of Missing Mechanisms in Dynamical Systems. *Interdisciplinary Math. Sci.*, 67-76, Vol. 8, 2010.
12. H. Crauel and F. Flandoli (1998), Additive noise destroys a pitchfork bifurcation, *Journal of Dynamics and Differential Equations* **10**, 259-274.
13. M. Davis, Pricing weather derivatives by marginal value, *Quantitative Finance* **1**(3)(2001), 305-308.
14. M. Denker, J. Duan and M. McCourt, Pseudorandom Numbers for Conformal Measures. *Dynamical Systems* **24** (2009), 439-457.
15. A. Du and J. Duan. A stochastic approach for parameterizing unresolved scales in a system with memory. *Journal of Algorithms & Computational Technology* **3**(2009), 393-405.
16. J. Duan, Stochastic Modeling of Unresolved Scales in Complex Systems. *Frontiers of Math. in China* **4** (2009), 425-436.
17. J. Duan, S. Luo and C. Wang (Eds.), *Recent Development in Stochastic Dynamics and Stochastic Analysis*. World Scientific, New Jersey, 2010.
18. J. Duan and B. Nadiga, Stochastic parameterization of large eddy simulation of geophysical flows. *Proc. American Math. Soc.* **135** (2007), 1187-1196.
19. J. Garcia-Ojalvo and J. M. Sancho, *Noise in Spatially Extended Systems*. Springer-Verlag, 1999.
20. C. W. Gardiner, *Handbook of Stochastic Methods*. Second Ed., Springer, New York,




Fig. 0.4. Squared Hellinger distance between $p(x, t)$ and the observation $q_2(x, t)$: $T = 5$ (top) and $T = 30$ (bottom).

- 1985.
21. V. Genon-Catalot and J. Jacod, On the estimation of the diffusion coefficient for multidimensional diffusion processes, *Ann. Inst. Henri Poincaré, Probabilités et Statistiques*. **29** (1993), 119-151.
22. V. Genon-Catalot and J. Jacod, On the estimation of the diffusion coefficient for diffusion processes, *J. Statist.* **21**(1994), 193-221.
23. V. Genon-Catalot, T. Jeantheau and C. Laredo, Parameter estimation for discretely observed stochastic volatility models, *Bernoulli* **5**(5) (1999), 855-872.
24. P. Hanggi, A. Alvarez-Chillida and M. Morillo Buzon (Eds.), New Horizons in Stochastic Complexity. Special Issue: *Physica A*, **351**(2005), 1-188.
25. K. Hasselmann, Stochastic climate models: Part I. Theory, *Tellus* **28** (1976), 473-485.
26. W. Horsthemke and R. Lefever, *Noise-Induced Transitions: Theory and Applications in Physics, Chemistry, and Biology*. Springer-Verlag, Berlin, second printing, 2007.
27. W. Huisinga, C. Schutte and A.M. Stuart, Extracting macroscopic stochastic dynamics: Model problems. *Comm. Pure Appl. Math.*, **56**2003, 234-269.
28. I. A. Ibragimov and R. Z. Has'minskii, *Statistical Estimation—Asymptotic Theory*. Springer, New York, 1981.
29. V. Isakov, *Inverse Problems for Partial Differential Equations*. Springer, New York, 1998.

30. J. Jacod, Parametric inference for discretely observed non-ergodic diffusions, *Bernoulli* **12**(3) (2006), 383-401.
31. W. Just, H. Kantz, C. Rodenbeck and M. Helm, Stochastic modelling: replacing fast degrees of freedom by noise. *J. Phys. A: Math. Gen.*, **34** (2001), 3199-3213.
32. F. C. Klebaner, *Introduction to Stochastic Calculus with Applications*. Imperial College Press, London, 2nd edition, 2005.
33. P. E. Kloeden and E. Platen, *Numerical solution of stochastic differential equations*, Springer-Verlag, 1992; second corrected printing, 1995.
34. Y. A. Kutoyants, *Statistical Inference for Ergodic Diffusion Processes*. Springer, New York, 2004.
35. C. E. Leith, Stochastic backscatter in a subgrid-scale model: Plane shear mixing layer. *Phys. Fluids A* **2** (1990), 297-299.
36. J. W.-B. Lin and J. D. Neelin, Considerations for stochastic convective parameterization, *J. Atmos. Sci.* 2002 Vol. 59, No. 5, pp. 959-975.
37. R. S. MacKay, Langevin equation for slow degrees of freedom of Hamiltonian systems, in "Nonlinear Dynamics and Chaos", eds M Theil, J Kurths, MC Romano, G Karolyi, A Moura (Springer, 2010) 89-102.
38. A. J. Majda, I. Timofeyev and E. Vanden Eijnden, Models for stochastic climate prediction, *Proc. NAS* **96** (1999), 14687-14691.
39. A. J. Majda, I. Timofeyev and E. Vanden Eijnden, A mathematical framework for stochastic climate models, *Comm. Pure Appl. Math.* **LIV** (2001), 891-974.
40. P. J. Mason and D. J. Thomson, Stochastic backscatter in large-eddy simulations of boundary layers. *J. Fluid Mech.* **242** (1992), 51-78.
41. F. Moss and P. V. E. McClintock (eds.), *Noise in Nonlinear Dynamical Systems*. Volume 1: Theory of Continuous Fokker-Planck Systems (2007); Volume 2: Theory of Noise Induced Processes in Special Applications (2009); Volume 3: Experiments and Simulations (2009). Cambridge University Press.
42. J. Nicolau, Introduction to the Estimation of Stochastic Differential Equations Based on Discrete Observations, Stochastic Finance, 2004 (Autumn School and International Conference).
43. B. Oksendal. *Stochastic differential equations*. Sixth Ed., Springer-Verlag, New York, 2003.
44. T. N. Palmer, G. J. Shutts, R. Hagedorn, F. J. Doblas-Reyes, T. Jung and M. Leutbecher. Representing model uncertainty in weather and climate prediction. *Annu. Rev. Earth Planet. Sci.* **33** (2005), 163-193.
45. C. Pasquero and E. Tziperman, Statistical parameterization of heterogeneous oceanic convection, *J. Phys. Oceanography*, **37** (2007), 214-229.
46. C. Penland and P. Sura, Sensitivity of an ocean model to "details" of stochastic forcing. In *Proc. ECMWF Workshop on Representation of Subscale Processes using Stochastic-Dynamic Models*. Reading, England, 6-8 June 2005.
47. N. D. Pearson and T. Sun, Exploiting the conditional density in estimating the term structure: an application to the Cox, Ingersoll and Ross model, *The Journal of Finance* **49**(4)(1994), 1279-1304.
48. B. L. S. Prakasa Rao, *Statistical Inference for Diffusion Type Processes*. London : Arnold ; New York : Oxford University Press, 1999.
49. B. L. Rozovskii, *Stochastic Evolution Equations*. Kluwer Academic Publishers, Boston, 1990.
50. P. Sagaut, *Large Eddy Simulation for Incompressible Flows*. Third Edition, Springer, 2005.
51. P. Sardeshmukh, Issues in stochastic parametrisation. In *Proc. ECMWF Workshop*

- on Represenation of Subscale Processes using Stochastic-Dynamic Models*. Reading, England, 6-8 June 2005.
52. U. Schumann, Stochastic backscatter of turbulent energy and scalar variance by random subgrid-scale fluxes. *Proc. R. Soc. Lond. A* **451** (1995), 293-318.
 53. N. G. Van Kampen, How do stochastic processes enter into physics? *Lecture Note in Phys.* **1250** (1987) 128-137.
 54. N. G. Van Kampen, *Stochastic Processes in Physics and Chemistry*, North-Holland, New York, 1981.
 55. W. M. Washington and C. L. Parkinson, *An Introduction to Three-Dimensional Climate Modeling*, Oxford Univ. Press, 1986.
 56. E. Waymire and J. Duan (Eds.). *Probability and Partial Differential Equations in Modern Applied Mathematics*. Springer-Verlag, 2005.
 57. D. S. Wilks, Effects of stochastic parameterizations in the Lorenz '96 system. *Q. J. R. Meteorol. Soc.* **131** (2005), 389-407.
 58. P. D. Williams. Modelling climate change: the role of unresolved processes. *Phil. Trans. R. Soc. A* (2005) **363**, 2931-2946.
 59. E. Wong and B. Hajek, *Stochastic Processes in Engineering Systems*, Spring-Verlag, New York, 1985.
 60. Y. Xu, R. Gu, H. Zhang, W. Xu and J. Duan, Stochastic Bifurcations in a Bistable Duffing-Van der Pol Oscillator with Colored Noise. *Phys. Rev. E.*, **83**, 056215 (2011).
 61. J. Yang and J. Duan, Quantifying Model Uncertainties in Complex Systems. *Progress in Prob.*, Vol. 65, p.49-80, Kohatsu-Higa, Arturo; Privault, Nicolas; Sheu, Shuenn-Jyi (Eds.), 2011. arxiv: 0912.0280

Quantifying Model Uncertainties in the Space of Probability Measures

Jinqiao Duan¹, Ting Gao¹ and Guowei He² *

*1. Institute for Pure and Applied Mathematics
University of California, Los Angeles, CA 90095, USA*

&

*Department of Applied Mathematics, Illinois Institute of Technology
Chicago, IL 60616, USA*

E-mail: duan@iit.edu tgao5@iit.edu

*2. Laboratory for Nonlinear Mechanics
Institute of Mechanics, Chinese Academy of Sciences
Beijing 100080, China*

E-mail: hgw@lnm.imech.ac.cn

Due to lack of scientific understanding, some mechanisms may be missing in mathematical modeling of complex phenomena in science and engineering. These mathematical models thus contain some uncertainties such as uncertain parameters. One method to estimate these parameters is based on pathwise observations, i.e., quantifying model uncertainty in the space of sample paths for system evolution. Another method is devised here to estimate uncertain parameters based on experimental observations of probability distributions for system evolution. This is called the quantification of model uncertainties in the space of probability measures. A few examples are presented to demonstrate this method, analytically or numerically.

Keywords: Model uncertainty; Stochastic differential equations (SDEs); Probability measures; Hellinger distance; Stationary probability density; Parameter estimation; Numerical simulation

2000 AMS Subject Classification : 60H40

1 . Introduction

In this chapter we discuss some issues about quantification of model uncertainties in complex dynamical systems.

*Partly supported by the NSF grant 1025422, NSFC grant 10971225, the open funding from the Laboratory for Nonlinear Mechanics at the Chinese Academy of Sciences, and the Fundamental Research Funds for the Central Universities, HUST 2010ZD037.

Mathematical models for scientific and engineering systems often involve with some uncertainties. We may roughly classify such uncertainties into two kinds. The first kind of uncertainties may be called *model uncertainty*. They are due to physical processes that are not well understood or not well-observed, and thus are not or not well represented in the mathematical models.

The second kind of uncertainties may be called *simulation uncertainty*. This arises in numerical simulations of multiscale systems that display a wide range of spatial and temporal scales, with no clear scale separation. Due to the limitations of computer power, not all scales of variability can be explicitly simulated or resolved.

These uncertainties are sometimes also called *unresolved scales*, as they are not represented (i.e., not *resolved*) in modeling or simulation. Although these unresolved scales may be very small or very fast, their long time impact on the resolved simulation may be delicate (i.e., may be negligible or may have significant effects, or in other words, uncertain). Thus, to take the effects of unresolved scales on the resolved scales into account, representations or parameterizations of these effects are desirable.

Model uncertainties have been considered in, for example,^{18,25,40,43–45} Research works relevant for parameterizing unresolved scales include,^{6–8,17,24,26,30,38,49,56,57} among others. Stochastically representing unresolved scales in fluid dynamics has considered as well.^{34,39,51}

In this chapter, we only consider model uncertainties. Specifically, we consider dynamical systems containing uncertain parameters and examine how to estimate these parameters, using observed probability distributions of the system evolution.

After briefly comment on estimating uncertain parameters based on observed sample paths for the system evolution in §2, we then, in §3, propose a method of estimating uncertain parameters based on observed probability distributions (i.e., probability measures) and present a few examples to demonstrate this method, analytically or numerically.

2 . Quantifying uncertainty in the space of paths

Since random fluctuations are common in the real world, mathematical models for complex systems are often subject to uncertainties, such as fluctuating forces, uncertain parameters, or random boundary conditions.^{18,25,40,52,55,58} Stochastic differential equations (SDEs) such as

$$dX = b(X)dt + \sigma(X)dB_t, \quad (2.1)$$

are appropriate models for many of these systems.^{5,48,55} Here B_t is a Brownian motion or Wiener process, the drift $b(X)$ and diffusion $\sigma(X)$ contain uncertain parameters, to be estimated based on observations.

For example, the Langevin type models are stochastic differential equations describing various phenomena in physics, biology, and other fields. SDEs are used to

model various price processes, exchange rates, and interest rates, among others, in finance. Noises in these SDEs may be modeled as a generalized time derivative of some distinguished stochastic processes, such as Brownian motion (BM) or other processes.

We are interested in estimating parameters contained in the stochastic differential equation (2.1), so that we obtain computational models useful for investigating complex dynamics under uncertainty.

Theoretical results on parameter estimations for SDEs driven by Brownian motion are relatively well developed,^{4,9,12,22,27,33,46} and various numerical simulations for these parameter estimations^{1,3,41,46} are also implemented. See⁶⁰ for a more recent review about estimating and computing uncertain parameters, when dynamical systems are submit to colored or non-Gaussian noises.

These research works on estimating uncertain parameters in dynamical systems are based on observations of sample paths. In the next section, we devise a method to estimate uncertain parameters based on observations of probability distributions of the system evolution.

3 . Quantifying uncertainty in the space of probability measures

Consider a dynamical system with model uncertainty, modeled by a scalar SDE

$$dX = b(X)dt + \sigma(X)dB_t, \quad X(0) = x_0, \quad (3.1)$$

where the drift $b(X)$ and diffusion $\sigma(X)$ contain uncertain parameters, to be estimated based on observations of probability distributions (i.e., probability measures) of the system paths X_t .

To this end, we need to introduce the Hellinger distance¹⁰ between two probability measures. It is used to quantify the similarity between two probability distributions.

This is a metric in the space of probability measures.

3 .1. Stationary case

Under appropriate conditions on b and σ (see,³¹ p.170.), such as, $b \leq 0$ and $\sigma \neq 0$ as well as some smoothness requirements, there exists a stationary probability density $p(x)$ for the SDE (3.1), as a solution of the steady Fokker-Planck equation,

$$p(x) = \frac{C}{\sigma^2(x)} e^{\int_{x_0}^x \frac{2b(y)}{\sigma^2(y)} dy},$$

where the positive normalization constant C is chosen so that $p \geq 0$ and $\int_{\mathbb{R}} p(x)dx = 1$, i.e.,

$$C \triangleq 1 / \int_{-\infty}^{\infty} \frac{e^{\int_{x_0}^x \frac{2b(y)}{\sigma^2(y)} dy}}{\sigma^2(x)} dx.$$

Note that x_0 here may be an arbitrary point so that the integral $\int_{x_0}^x \frac{2b(y)}{\sigma^2(y)} dy$ exists (say, take $x_0 = 0$ if that is convenient).

Example 3 .1. (a) A special case: Langevin equation

$$dX = -bXdt + dB_t,$$

with parameter $b > 0$. Given an “observation” of the stationary probability density $q(x) = \frac{1}{\sqrt{\pi}}e^{-x^2}$. Find a b so that the Hellinger distance $F(b) = \frac{1}{2} \int_{\mathbb{R}} [\sqrt{p(x)} - \sqrt{q(x)}]^2 dx$ is minimized.

Hint: Insert p, q into the Hellinger distance. $F(b)$ is an algebraic function of parameter $b > 0$ and thus we use deterministic calculus to find a minimizer b (possibly by hand, or Matlab if needed). Note: $\int_{\mathbb{R}} e^{-z^2} dz = \sqrt{\pi}$.

(b) A more general case:

$$dX = b(X)dt + dB_t,$$

with function $b(x) \leq 0$. Given an “observation” of the stationary probability density $q(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ (the Cauchy distribution). Find a function $b(x) \leq 0$ so that the Hellinger distance $F(b(x)) = \frac{1}{2} \int_{\mathbb{R}} [\sqrt{p(x)} - \sqrt{q(x)}]^2 dx$ is minimized.

Hint: Insert p, q into the Hellinger distance. $F(b(x))$ is a functional of $b(x)$ and thus we use calculus of variations (on $F(b(x))$) to find a minimizer b . Derive the Euler-Lagrange equation to be satisfied by $b(x)$, together with appropriate boundary conditions (needed for $p \geq 0$ and $\int_{\mathbb{R}} p dx = 1$). Can you devise an algorithm to simulate for $b(x)$?

Solution:

(a) From the Fokker-Planck equation of

$$dX_t = -bX_t dt + dB_t, \quad X_0 = 0,$$

we have the stationary probability density

$$P(x) = \frac{\sqrt{b}}{\sqrt{\pi}} e^{-bx^2}.$$

To minimize the Hellinger distance $F(b)$, we need

$$\begin{aligned} F'(b) &= \frac{1}{2} \int_{\mathbb{R}} e^{-bx^2} \left(\frac{1}{2\sqrt{\pi b}} - \frac{\sqrt{b}}{\sqrt{\pi}} x^2 \right) dx - \frac{1}{\sqrt{\pi}} \int_{\mathbb{R}} e^{-\frac{(1+b)x^2}{2}} \left(\frac{1}{4} b^{-\frac{3}{4}} - \frac{x^2}{2} b^{\frac{1}{4}} \right) dx \\ &= \frac{1}{\sqrt{2}} b^{\frac{1}{4}} (1+b)^{-\frac{3}{2}} - \frac{1}{2\sqrt{2}} b^{-\frac{3}{4}} (1+b)^{-\frac{1}{2}} = 0. \end{aligned}$$

Therefore,

$$b^{\frac{1}{4}} (1+b)^{-\frac{3}{2}} = \frac{b^{-\frac{3}{4}} (1+b)^{-\frac{1}{2}}}{2} \Rightarrow b = 1.$$

(b) From the Fokker-Planck equation of

$$dX_t = b(X_t)dt + dB_t, \quad X_0 = 0,$$

we could have the stationary probability density

$$P(x) = \frac{e^{2 \int_0^x b(y)dy}}{\int_{-\infty}^{\infty} e^{2 \int_0^x b(y)dy} dx}.$$

Using calculus of variation, we need

$$\begin{aligned} & F(b(x) + \varepsilon h(x)) \\ &= \frac{1}{2} \int_{\mathbb{R}} \left[\frac{e^{2 \int_0^x (b(y) + \varepsilon h(y))dy}}{\int_{-\infty}^{\infty} e^{2 \int_0^x (b(y) + \varepsilon h(y))dy} dx} - \frac{2e^{\int_0^x (b(y) + \varepsilon h(y))dx}}{\sqrt{\pi} \sqrt{1+x^2} \sqrt{\int_{-\infty}^{\infty} e^{2 \int_0^x (b(y) + \varepsilon h(y))dy} dx}} + \frac{1}{\pi(1+x^2)} \right] dx \\ &= 0. \end{aligned}$$

This is equivalent to

$$\int_{\mathbb{R}} e^{\int_0^x b(w)dw} \left(e^{\int_0^x b(w)dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx}}{\sqrt{\pi(1+x^2)}} \right) \cdot \int_{\mathbb{R}} \left(e^{2 \int_0^z b(w)dw} \int_z^x h(y)dy \right) dz = 0.$$

When changing the order of integration (first on y and z then on y and x), we have

$$\left\{ \int_y^{\infty} e^{\int_0^x b(w)dw} \left(e^{\int_0^x b(w)dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx}}{\sqrt{\pi(1+x^2)}} \right) dx \cdot \int_{-\infty}^y e^{2 \int_0^z b(w)dw} dz \right\} h(y)dy = 0$$

holds for all $h(y)$.

Therefore,

$$\int_y^{\infty} e^{\int_0^x b(w)dw} \left(e^{\int_0^x b(w)dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx}}{\sqrt{\pi(1+x^2)}} \right) dx \cdot \int_{-\infty}^y e^{2 \int_0^z b(w)dw} dz = 0.$$

And since $\int_{-\infty}^y e^{2 \int_0^z b(w)dw} dz > 0$, we have

$$\int_y^{\infty} e^{\int_0^x b(w)dw} \left(e^{\int_0^x b(w)dw} - \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx}}{\sqrt{\pi(1+x^2)}} \right) dx = 0$$

for $y \in \mathbb{R}$. When taking the derivative of the above equation with respect to y , we have

$$e^{\int_0^y b(w)dw} = \frac{\sqrt{\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx}}{\sqrt{\pi(1+y^2)}}, \quad \forall y \in \mathbb{R}.$$

Therefore, after taking 'square' and 'ln' on both sides of the above equation, we have

$$2 \int_0^y b(w)dw = \ln\left(\int_{\mathbb{R}} e^{2 \int_0^x b(w)dw} dx\right) - \ln(\pi(1+y^2)).$$

Taking derivative with respect to y , we have

$$b(y) = -\frac{y}{\pi(1+y^2)} \quad y \in \mathbb{R}.$$

only need $\int_0^y b(w)dw < 0$ **for all** $y \in \mathbb{R}$.

3.2. Time-dependent case

Consider a scalar SDE

$$dX = b(X)dt + \sigma(X)dB_t, \quad X(0) = x. \quad (3.2)$$

The Fokker-Planck equation

$$p_t = \frac{1}{2}(\sigma^2(x)p(x,t))_{xx} - (b(x)p(x,t))_x, \quad (3.3)$$

28

Example 3.2. Consider a scalar SDE

$$dX = -b \sin(X)dt + \sqrt{2} dB_t, \quad X(0) = 0. \quad (3.4)$$

(1) Assume that an observation obtained for p to be

$$q_1(x,t) = \frac{1}{2\sqrt{\pi t}} e^{-\frac{x^2}{4t}}. \quad (3.5)$$

(2) Assume that another observation obtained for p to be

$$q_2(x,t) = \frac{\sqrt{t}}{\pi(t+x^2)}. \quad (3.6)$$

The Fokker-Planck equation for (3.4) is

$$p_t = p_{xx} + (b \sin(x)p)_x. \quad (3.7)$$

Find b based on the above two observation.

Solution: In this case, we define the Hellinger distance:

$$H_i(b) = \max_{t \in [0, T]} \int_{-\infty}^{\infty} (\sqrt{q_i(x,t)} - \sqrt{p(x,t,b)})^2 dx$$

where $i = 1, 2$. We need to calculate b by minimizing $H_i(b)$.

Now it's not easy to get the analytical solution of $p(x, t)$, so we do some computer simulations instead. Assuming that the initial distribution $p(x, 0)$ is a delta function δ_0 on $(-\infty, \infty)$, we can solve the above Fokker-Planck equation numerically, the solution is in the following figures.




Fig. 0.1. Solutions of Fokker-Planck equation when $b = 1$. In the plot, $t = 5$ (left) and $t = 30$ (right).

For the observation $q_1(x, t)$, it's the solution of $p_t = p_{xx}$ with delta function δ_0 as the initial condition. Hence, the corresponding SDE of $q_1(x, t)$ is $dX = \sqrt{2} dB_t$, $X(0) = 0$. We have its distribution shown below:




Fig. 0.2. Observation distribution $q_1(x, t)$. In the plot, $t = 5$ (left) and $t = 30$ (right).

By the definition $H_1(b) = \max_{t \in [0, T]} \int_{-\infty}^{\infty} (\sqrt{q_1(x, t)} - \sqrt{p(x, t, b)})^2 dx$, we have the plot of $H_1(b)$ in the following figures. And whatever T is, $H_1(b)$ is always minimized when $b = 0$.

For the observation $q_2(x, t)$, we have its distribution like

By the definition $H_2(b) = \max_{t \in [0, T]} \int_{-\infty}^{\infty} (\sqrt{q_2(x, t)} - \sqrt{p(x, t, b)})^2 dx$, we have the plot of $H_2(b)$ in the following figures. So we can see that if $T = 5$, $H_2(b)$ is minimized when $b = 0.7$ and if $T = 30$, $H_2(b)$ is minimized when $b = 0.6$.

Acknowledgements. We would like to thank Huijie Qiao and Xiangjun Wang for helpful discussions.

References

1. Y. Aït-Sahalia (2002), Maximum-likelihood estimation of discretely-sampled diffusions: a closed-form approximation approach, *Econometrica* **70**, 223-262.
2. Y. Aït-Sahalia and P. A. Mykland (2004), Estimators of diffusions with randomly




Fig. 0.3. Hellinger distance between pdf $p(x, t)$ and observation $q_1(x, t)$. In the plot, $T = 5$ (left) and $T = 30$ (right).

- spaced discrete observations: a general theory, *The Annals of Statistics* **32**(5), 2186-2222
3. Y. Aït-Sahalia and P. A. Mykland (2003), The effects of random and discrete sampling when estimating continuous-time diffusions, *Econometrica* **71**(2), 483-549.
 4. S. Alizadeh, M. W. Brandt and F. X. Diebold (2002), Range-based estimation of stochastic volatility models, *The Journal of Finance* **57**(3), 1047-1091.
 5. L. Arnold. *Random Dynamical Systems*. Springer, New York, 1998.
 6. L. Arnold, Hasselmann's program revisited: The analysis of stochasticity in deterministic climate models. In J.-S. von Storch and P. Imkeller, editors, *Stochastic climate models*. pages 141-158, Boston, 2001. Birkhäuser.
 7. P. S. Berloff, Random-forcing model of the mesoscale oceanic eddies. *J. Fluid Mech.* **529** (2005), 71-95.
 8. L.C. Berselli, T. Iliescu and W. J. Layton. *Mathematics of Large Eddy Simulation of Turbulent Flows*. Springer Verlag, 2005.
 9. J. P. N. Bishwal, *Parameter Estimation in Stochastic Differential Equations*, Springer, New York, 2007.
 10. S. Cha, Comprehensive Survey on Distance/Similarity Measures between Probability Density Functions. *International Journal of Mathematical Models and Methods in Applied Sciences*, vol. 1(4), 2007 pp.300-307.




Fig. 0.4. Observation distribution $q_2(x, t)$. In the plot, $t = 5$ (left) and $t = 30$ (right).

11. B. Chen and J. Duan, Stochastic Quantification of Missing Mechanisms in Dynamical Systems. *Interdisciplinary Math. Sci.*, 67-76, Vol. 8, 2010.
12. M. Davis (2001), Pricing weather derivatives by marginal value, *Quantitative Finance* **1**(3), 305-308.
13. M. Denker, J. Duan and M. McCourt, Pseudorandom Numbers for Conformal Measures. *Dynamical Systems* **24** (2009), 439-457.
14. A. Du and J. Duan. A stochastic approach for parameterizing unresolved scales in a system with memory. *Journal of Algorithms & Computational Technology* **3**(2009), 393-405.
15. J. Duan, Stochastic Modeling of Unresolved Scales in Complex Systems. *Frontiers of Math. in China* **4** (2009), 425-436.
16. J. Duan, S. Luo and C. Wang (Eds.), *Recent Development in Stochastic Dynamics and Stochastic Analysis*. World Scientific, New Jersey, 2010.
17. J. Duan and B. Nadiga, Stochastic parameterization of large eddy simulation of geophysical flows. *Proc. American Math. Soc.* **135** (2007), 1187-1196.
18. J. Garcia-Ojalvo and J. M. Sancho, *Noise in Spatially Extended Systems*. Springer-Verlag, 1999.
19. C. W. Gardiner, *Handbook of Stochastic Methods*. Second Ed., Springer, New York, 1985.
20. V. Genon-Catalot and J. Jacod (1993), On the estimation of the diffusion coefficient




Fig. 0.5. Hellinger distance between pdf $p(x, t)$ and observation $q_2(x, t)$. In the plot, $T = 5$ (left) and $T = 30$ (right).

- for multidimensional diffusion processes, *Ann. Inst. Henri Poincaré, Probabilités et Statistiques*. **29**, 119-151.
21. V. Genon-Catalot and J. Jacod (1994), On the estimation of the diffusion coefficient for diffusion processes, *J. Statist.* **21**, 193-221.
 22. V. Genon-Catalot, T. Jeantheau and C. Laredo (1999), Parameter estimation for discretely observed stochastic volatility models, *Bernoulli* **5**(5), 855-872.
 23. P. Hanggi, A. Alvarez-Chillida and M. Morillo Buzon (Eds.), New Horizons in Stochastic Complexity. Special Issue: *Physica A*, **351**(2005), 1-188.
 24. K. Hasselmann, Stochastic climate models: Part I. Theory, *Tellus* **28** (1976), 473-485.
 25. W. Horsthemke and R. Lefever, *Noise-Induced Transitions: Theory and Applications in Physics, Chemistry, and Biology*. Springer-Verlag, Berlin, second printing, 2007.
 26. W. Huisinga, C. Schutte and A.M. Stuart, Extracting macroscopic stochastic dynamics: Model problems. *Comm. Pure Appl. Math.*, **56**2003, 234-269.
 27. I. A. Ibragimov and R. Z. Has'minskii, *Statistical Estimation—Asymptotic Theory*. Springer, New York, 1981.
 28. V. Isakov, *Inverse Problems for Partial Differential Equations*. Springer, New York, 1998.
 29. J. Jacod (2006), Parametric inference for discretely observed non-ergodic diffusions, *Bernoulli* **12**(3), 383-401.

30. W. Just, H. Kantz, C. Rodenbeck and M. Helm, Stochastic modelling: replacing fast degrees of freedom by noise. *J. Phys. A: Math. Gen.*, **34** (2001), 3199–3213.
31. F. C. Klebaner, *Introduction to Stochastic Calculus with Applications*. Imperial College Press, London, 2nd edition, 2005.
32. P. E. Kloeden and E. Platen, *Numerical solution of stochastic differential equations*, Springer-Verlag, 1992; second corrected printing, 1995.
33. Y. A. Kutoyants, *Statistical Inference for Ergodic Diffusion Processes*. Springer, New York, 2004.
34. C. E. Leith, Stochastic backscatter in a subgrid-scale model: Plane shear mixing layer. *Phys. Fluids A* **2** (1990), 297–299.
35. J. W.-B. Lin and J. D. Neelin, Considerations for stochastic convective parameterization, *J. Atmos. Sci.* 2002 Vol. 59, No. 5, pp. 959–975.
36. R. S. MacKay, Langevin equation for slow degrees of freedom of Hamiltonian systems, in “*Nonlinear Dynamics and Chaos*”, eds M Theil, J Kurths, MC Romano, G Karolyi, A Moura (Springer, 2010) 89–102.
37. A. J. Majda, I. Timofeyev and E. Vanden Eijnden, Models for stochastic climate prediction, *Proc. NAS* **96** (1999), 14687–14691.
38. A. J. Majda, I. Timofeyev and E. Vanden Eijnden, A mathematical framework for stochastic climate models, *Comm. Pure Appl. Math.* **LIV** (2001), 891–974.
39. P. J. Mason and D. J. Thomson, Stochastic backscatter in large-eddy simulations of boundary layers. *J. Fluid Mech.* **242** (1992), 51–78.
40. F. Moss and P. V. E. McClintock (eds.), *Noise in Nonlinear Dynamical Systems*. Volume 1: Theory of Continuous Fokker-Planck Systems (2007); Volume 2: Theory of Noise Induced Processes in Special Applications (2009); Volume 3: Experiments and Simulations (2009). Cambridge University Press.
41. J. Nicolau (2004), Introduction to the Estimation of Stochastic Differential Equations Based on Discrete Observations, Stochastic Finance 2004 (Autumn School and International Conference).
42. B. Oksendal. *Stochastic differential equations*. Sixth Ed., Springer-Verlag, New York, 2003.
43. T. N. Palmer, G. J. Shutts, R. Hagedorn, F. J. Doblas-Reyes, T. Jung and M. Leutbecher. Representing model uncertainty in weather and climate prediction. *Annu. Rev. Earth Planet. Sci.* **33** (2005), 163–193.
44. C. Pasquero and E. Tziperman, Statistical parameterization of heterogeneous oceanic convection, *J. Phys. Oceanography*, **37** (2007), 214–229.
45. C. Penland and P. Sura, Sensitivity of an ocean model to “details” of stochastic forcing. In *Proc. ECMWF Workshop on Representation of Subscale Processes using Stochastic-Dynamic Models*. Reading, England, 6–8 June 2005.
46. N. D. Pearson and T. Sun (1994), Exploiting the conditional density in estimating the term structure: an application to the Cox, Ingersoll and Ross model, *The Journal of Finance* **49**(4), 1279–1304.
47. B.L.S. Prakasa Rao. *Statistical Inference for Diffusion Type Processes*. London : Arnold ; New York : Oxford University Press, 1999.
48. B. L. Rozovskii, *Stochastic Evolution Equations*. Kluwer Academic Publishers, Boston, 1990.
49. P. Sagaut, *Large Eddy Simulation for Incompressible Flows*. Third Edition, Springer, 2005.
50. P. Sardeshmukh, Issues in stochastic parametrisation. In *Proc. ECMWF Workshop on Representation of Subscale Processes using Stochastic-Dynamic Models*. Reading, England, 6–8 June 2005.

51. U. Schumann, Stochastic backscatter of turbulent energy and scalar variance by random subgrid-scale fluxes. *Proc. R. Soc. Lond. A* **451** (1995), 293-318.
52. N. G. Van Kampen, How do stochastic processes enter into physics? *Lecture Note in Phys.* **1250** (1987) 128-137.
53. N. G. Van Kampen (1981), *Stochastic Processes in Physics and Chemistry*, North-Holland, New York.
54. W. M. Washington and C. L. Parkinson, *An Introduction to Three-Dimensional Climate Modeling*, Oxford Univ. Press, 1986.
55. E. Waymire and J. Duan (Eds.). *Probability and Partial Differential Equations in Modern Applied Mathematics*. Springer-Verlag, 2005.
56. D. S. Wilks, Effects of stochastic parameterizations in the Lorenz '96 system. *Q. J. R. Meteorol. Soc.* **131** (2005), 389-407.
57. P. D. Williams. Modelling climate change: the role of unresolved processes. *Phil. Trans. R. Soc. A* (2005) **363**, 2931-2946.
58. E. Wong and B. Hajek (1985), *Stochastic Processes in Engineering Systems*, Springer-Verlag, New York.
59. Y. Xu, R. Gu, H. Zhang, W. Xu and J. Duan, Stochastic Bifurcations in a Bistable Duffing-Van der Pol Oscillator with Colored Noise. *Phys. Rev. E.*, **83**, 056215 (2011).
60. J. Yang and J. Duan, Quantifying Model Uncertainties in Complex Systems. *Progress in Prob.*, Vol. 65, p.49-80, Kohatsu-Higa, Arturo; Privault, Nicolas; Sheu, Shuenn-Jyi (Eds.), 2011. arxiv: 0912.0280